

## Pocket CONCUSSION RECOGNITION TOOL™

To help identify concussion in children, youth and adults



### RECOGNIZE & REMOVE

Concussion should be suspected **if one or more** of the following visible clues, signs, symptoms or errors in memory questions are present.

#### 1. Visible clues of suspected concussion

Any one or more of the following visual clues can indicate a possible concussion:

- Loss of consciousness or responsiveness
- Lying motionless on ground/Slow to get up
- Unsteady on feet / Balance problems or falling over/Incoordination
- Grabbing/Clutching of head
- Dazed, blank or vacant look
- Confused/Not aware of plays or events

#### 2. Signs and symptoms of suspected concussion

Presence of any one or more of the following signs & symptoms may suggest a concussion:

- |                          |                            |
|--------------------------|----------------------------|
| - Loss of consciousness  | - Headache                 |
| - Seizure or convulsion  | - Dizziness                |
| - Balance problems       | - Confusion                |
| - Nausea or vomiting     | - Feeling slowed down      |
| - Drowsiness             | - "Pressure in head"       |
| - More emotional         | - Blurred vision           |
| - Irritability           | - Sensitivity to light     |
| - Sadness                | - Amnesia                  |
| - Fatigue or low energy  | - Feeling like "in a fog"  |
| - Nervous or anxious     | - Neck Pain                |
| - "Don't feel right"     | - Sensitivity to noise     |
| - Difficulty remembering | - Difficulty concentrating |

© 2013 Concussion in Sport Group

### 3. Memory function

Failure to answer any of these questions correctly may suggest a concussion.

- "What venue are we at today?"
- "Which half is it now?"
- "Who scored last in this game?"
- "What team did you play last week / game?"
- "Did your team win the last game?"

**Any athlete with a suspected concussion should be IMMEDIATELY REMOVED FROM PLAY, and should not be returned to activity until they are assessed medically. Athletes with a suspected concussion should not be left alone and should not drive a motor vehicle.**

It is recommended that, in all cases of suspected concussion, the player is referred to a medical professional for diagnosis and guidance as well as return to play decisions, even if the symptoms resolve.

#### RED FLAGS

**If ANY of the following are reported then the player should be safely and immediately removed from the field. If no qualified medical professional is available, consider transporting by ambulance for urgent medical assessment:**

- |  |                                 |
|--|---------------------------------|
| - Athlete complains of neck pain               | - Deteriorating conscious state |
| - Increasing confusion or irritability         | - Severe or increasing headache |
| - Repeated vomiting                            | - Unusual behaviour change      |
| - Seizure or convulsion                        | - Double vision                 |
| - Weakness or tingling/burning in arms or legs |                                 |

#### Remember:

- In all cases, the basic principles of first aid (danger, response, airway, breathing, circulation) should be followed.
- Do not attempt to move the player (other than required for airway support) unless trained to do so
- Do not remove helmet (if present) unless trained to do so.

from McCrory et. al, Consensus Statement on Concussion in Sport. Br J Sports Med 47 (5), 2013

© 2013 Concussion in Sport Group



## Pocket CRT

*Br J Sports Med* 2013 47: 267

---

Updated information and services can be found at:  
<http://bjsm.bmj.com/content/47/5/267.citation>

---

*These include:*

**Email alerting  
service**

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>